

Letter to the Editor

Hepatitis A - the neglected sexually transmissible disease

Mark J Ferson, Director, South Eastern Sydney Public Health Unit, Zetland, New South Wales, Locked Bag 88, Randwick NSW 2031

To the Editor: Dore and Kaldor provide valuable guidance on the directions a coordinated national system for surveillance of sexually transmissible diseases (STD) might take.¹ They did not attempt to provide an outline of what diseases might fall into this group, or what data should be collected for each disease, although a recent review of a decade of STD surveillance in Victoria examined data on syphilis, gonorrhoea, chlamydia, genital herpes, genital warts, hepatitis B, chancroid, lymphogranuloma venereum and donovanosis.²

However, it would seem timely to draw attention again to hepatitis A as a sexually transmissible disease, at least in some urban populations. The epicentre of a prolonged epidemic of hepatitis A in 1991-92 among homosexual men, was the gay community of the inner Sydney suburbs. A crude incidence of 136 cases per 100,000 population was recorded for 1991 in eastern Sydney, compared to 13

per 100,000 for Australia as a whole.³ The same epidemic was observed in Melbourne⁴ and simultaneously in other cities around the world.⁵ A similar, though smaller, epidemic occurred in 1995-96 in Sydney.

At the onset of the 1991-92 epidemic, the then Eastern Sydney Public Health Unit established a standalone hepatitis A database where risk factor information, in particular sexual preference, was kept in de-identified format. Maintenance of the database through subsequent epidemics in other risk groups^{6,7} has been invaluable in helping direct prevention activities, including education regarding hygiene and appropriate use of normal human immunoglobulin and hepatitis A vaccine. Analysis of the data revealed that during the 1991-92 and 1995-96 epidemics, adult males comprised almost 90% of all cases. This led to a peak incidence among young males, approaching 500 cases per 100,000 (or 0.5%) per year. Information on sexual preference was recorded in 75% of adult males, and of these, men who have sex with men contributed 80-90% of cases. In Australian cities, hepatitis A appears to behave as a sexually transmissible disease, and its proper surveillance requires collection of comprehensive risk factor data including sexual preference.

References

1. Dore GJ, Kaldor JM. Sexually transmissible diseases surveillance in Australia: towards a coordinated national system. *Comm Dis Intell* 1998;22:49-52.
2. Crofts N, Gertig DM, Stevenson E, et al. Surveillance for sexually transmissible diseases in Victoria, 1983 to 1992. *Aust J Public Health* 1994;18:433-9.
3. Stokes M-L, Ferson MJ, Young LC. Outbreak of hepatitis A among homosexual men in Sydney. *Am J Public Health* 1997; 87:-39-41.
4. Stewart T, Crofts N. An outbreak of hepatitis A among homosexual men in Melbourne. *Med J Aust* 1993;158:519-21.
5. CDC. Hepatitis A among homosexual men - United States, Canada and Australia. *MMWR* 1992; 41:155,161-4.
6. Ferson MJ, Young LC, Robertson PW, Whybin LR. Hepatitis A outbreak in a preschool in Eastern Sydney. *Comm Dis Intell* 1994;18:82-83.
7. Ferson MJ, Young LC. Preliminary report - Hepatitis A in injecting drug users. *Comm Dis Intell* 1994;18:655.

Notice to readers

Master of Applied Epidemiology - scholarships

The Master of Applied Epidemiology is a challenging, hands-on program which develops high level expertise in applying epidemiological methods to surveillance, outbreak investigation and disease control. The program is a leading member of the global Network of Training Programs in Epidemiology and Public Health Interventions that is collaborating with WHO's training and emergency response activities.

It is predominantly field-based, involving 21 months of field placement in a public health department or institution, and 12 weeks of intensive coursework (in four blocks) in Canberra.

Graduates of the medical and health sciences, with at least two years' health-related work experience, are eligible to apply for the two-year, full-time scholarships. Selection will be on the basis of academic merit and work experience. Scholars will receive a tax-free stipend of \$32,032 pa.

Applications close: 10 July 1998.

For further information and application papers, please contact:

Ros Hales, MAE Administrator, NCEPH, The Australian National University, Canberra ACT 0200

Telephone: (02) 6249 2790

Facsimile: (02) 6249 0740

Email: Ros.Hales@anu.edu.au

<http://www-nceph.anu.edu.au/student.htm>