

Update on the outbreak of cholera affecting Pohnpei State in the Federated States of Micronesia

Adapted from a report forwarded from the Pacific Public Health Surveillance Network¹

The outbreak was first recognised on 17 April 2000. As of 26 June 1,596 cases of watery diarrhoea have been reported – a figure based on inpatient and outpatient records from the Pohnpei Hospital together with data provided by community-based dispensaries. Nine deaths have been associated with the outbreak. *Vibrio cholerae* has been isolated by the Pohnpei Hospital and confirmed as the Ogawa serotype. Of these suspected cholera cases, 954 meet the World Health Organization case definition for cholera.² The epidemic is consistent with an initial point-source outbreak at a large funeral at Kitti, followed by a mixture of person-to-person transmission and additional point source outbreaks in a clockwise direction around the island.

Up to 27 June 2000, 368 (92%) of the 399 admissions to hospital, and 625 (80%) of 777 outpatient and emergency department presentations with acute watery diarrhoea, met the WHO case definition for cholera. Of the hospitalised cases, 40% were male, with adults more commonly infected. There appeared to be a bimodal distribution, with highest rates in males in the 65-74 year age group, and in females 20-24 years or over 75 years old.

Of the 777 non-hospitalised suspected cholera patients, 50% were male and 119 were less than five years of age. Due to possible multiple outpatient visits by the same patient, together with recording and data entry errors, the figures for non-hospitalised cases need to be treated with caution.

1. Information provided by the Federated States of Micronesia Department of Health via the Pacific Public Health Surveillance Network, 30 June 2000.
2. For surveillance purposes, in an epidemic situation, the case definition for suspected cholera is a person five years of age or greater developing acute watery diarrhoea.

Overseas briefs

Source: World Health Organization (WHO)

This material has been summarised from information on the WHO Internet site. A link to this site can be found under 'Other Australian and international communicable diseases sites' on the CDI homepage.

Accidental exposure to smallpox vaccine in the Russian Federation

The recent report of illness amongst eight young children in Vladivostok who had played with discarded ampoules of smallpox vaccine, has now been confirmed by the Ministry of Health of the Russian Federation. Laboratory confirmation of the illness in the children is being sought. The report has evoked much public concern. In some of the reports, there were misconceptions about the components of the vaccine used to prevent smallpox, and about why any country might still be retaining stocks of smallpox vaccine. This note aims to clarify these issues.

1) Smallpox vaccine is not made from smallpox virus.

The vaccine which was used for centuries to vaccinate against smallpox was not made from smallpox, but from vaccinia virus. Vaccinia is a different virus from the virus that causes smallpox. However, it is a member of the same family of viruses to which the smallpox virus belongs. The smallpox virus is also known as variola virus. Mass vaccinations with smallpox vaccine made from vaccinia virus led to the eradication of smallpox announced by WHO in 1980. People vaccinated with smallpox vaccine (vaccinia) develop reactions to it which range from mild and transient to severe and, very rarely, fatal.

2) Two countries still keep smallpox virus (variola) stocks.

Although smallpox disease has been eradicated, two laboratories still hold stocks of smallpox virus (variola). These are the WHO Collaborating Centres in Atlanta (USA) and Koltsovo (Russian Federation).

3) Many countries still hold smallpox vaccine (vaccinia) stocks.

WHO recommends that countries which still have stocks of smallpox vaccine (vaccinia) maintain these stocks. This recommendation has been made for two reasons. Firstly, small amounts of vaccine are still needed to vaccinate laboratory personnel handling vaccinia virus and other members of this virus family. Some of these viruses are found in nature and cause illness among animals, and some are used in research to make new, safer vaccines against a variety of infectious diseases. Secondly, smallpox vaccine (vaccinia) will also be needed in case of a deliberate or accidental release of smallpox virus (variola), which is a very unlikely event but currently of great concern to some countries.

4) Disposal of biological materials and pharmaceuticals

All biological materials and pharmaceuticals such as vaccines, drugs and diagnostic specimens should be disposed of safely. Some may require inactivation before disposal. This can be accomplished by autoclaving or incineration.

Acute haemorrhagic fever syndrome in Afghanistan

An outbreak of acute haemorrhagic fever syndrome has been reported from an isolated village in Gulran district, Afghanistan. Disease symptoms are compatible with Crimean-Congo haemorrhagic fever (CCHF). An international team, coordinated by WHO, arrived in the affected area on 16 June to control and investigate this outbreak. The team comprises experts from the WHO Collaborating Centre at the National Institute of Virology (NIV - South Africa), Epicentre (France) and WHO. Preliminary findings indicate that cases began at the beginning of May and are continuing to occur. Twenty-five suspect cases, including 15 deaths, were identified by the team. Samples were collected and transported to NIV where diagnostic laboratory tests will be performed to establish the aetiology of the outbreak.

The team has provided basic protective materials (gloves, masks), disinfectants (chlorine bleach) and instructions for

their use in caring for patients with bleeding symptoms. An isolation area has been identified in the hospital in Herat to deal with haemorrhagic patients and will be equipped by WHO. Training in barrier nursing will be provided to medical and nursing staff.

E. coli O157 in Canada

The Public Health Unit for the town of Walkerton, Ontario (population 5,000) has reported an outbreak of *E. coli* O157. As at 30 May, 5 people had died and 27 were hospitalised. Exposure to *E. coli* is believed to have occurred between 12-15 May. Tests have confirmed *E. coli* contamination in the Walkerton water supply system. On 21 May, the Public Health Unit issued a "Boil Water Order" to the residents of Walkerton which is still in effect. Investigations are under way by provincial and federal health and environmental authorities.

Editor: Angela Merianos **Associate Editor:** Jenny Thomson

Deputy Editor: Ian Griffith

Editorial and Production Staff

Alison Milton, Gail Bird, Margo Eyeson-Annan, Ming Lin

Editorial Advisory Board

Charles Watson (Chair), Mary Beers, Margaret Burgess, Scott Cameron, John Kaldor, Margery Kennett, Cathy Mead

Subscriptions

CanPrint, PO Box 7456, Canberra Mail Centre, ACT, 2610;
Fax: +61 2 6295 4888 (Overseas) or (02) 6295 4888 (Australia).

Website

<http://www.health.gov.au/pubhlth/cdi/cdihtml.htm>

Contributions

Contributions covering any aspects of communicable diseases are invited. All contributions are subject to the normal refereeing process.

Instructions to authors can be found in *CDI* 2000;24:5-6

Copyright

© Commonwealth of Australia 2000

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968* no part may be reproduced by any process without prior written permission from the Commonwealth available from AusInfo. Requests and inquires concerning reproduction and rights should be addressed to the Manager, Legislative Services, AusInfo, GPO Box 1920, Canberra ACT 2601.

Contacts other than subscriptions

CDI is produced every four weeks by the National Centre for Disease Control, Department of Health and Aged Care, GPO Box 9848, Canberra, ACT, 2601; Fax: (02) 6289 7791, Phone: (02) 6289 8245; email: cdi.editor@health.gov.au.

This journal is indexed by *Index Medicus* and Medline.

Opinions expressed in *CDI* are those of the authors and not necessarily those of the Department of Health and Aged Care or the Communicable Diseases Network Australia New Zealand. Data may be subject to revision.